

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/435, 14/715, 16/28, C12N 15/12, 15/19</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/35986</b> <b>(43) International Publication Date:</b> 20 August 1998 (20.08.98)															
<b>(21) International Application Number:</b> PCT/US98/02239 <b>(22) International Filing Date:</b> 11 February 1998 (11.02.98)  <b>(30) Priority Data:</b> <table border="0" style="width: 100%;"><tr><td style="width: 30%;">08/799,861</td><td style="width: 40%;">13 February 1997 (13.02.97)</td><td style="width: 30%;">US</td></tr><tr><td>08/815,255</td><td>12 March 1997 (12.03.97)</td><td>US</td></tr><tr><td>08/829,536</td><td>28 March 1997 (28.03.97)</td><td>US</td></tr><tr><td>08/869,852</td><td>4 June 1997 (04.06.97)</td><td>US</td></tr><tr><td>08/883,036</td><td>26 June 1997 (26.06.97)</td><td>US</td></tr></table> <b>(71) Applicant:</b> IMMUNEX CORPORATION [US/US]; Law Dept., 51 University Street, Seattle, WA 98101 (US).  <b>(72) Inventors:</b> RAUCH, Charles; 6455 N.E. Dapple Court, Bainbridge Island, WA 98110 (US). WALCZAK, Henning; 101 West Olympic Place, Seattle, WA 98119 (US).  <b>(74) Agent:</b> ANDERSON, Kathryn, A.; Immunex Corporation, Law Dept., 51 University Street, Seattle, WA 98101 (US).		08/799,861	13 February 1997 (13.02.97)	US	08/815,255	12 March 1997 (12.03.97)	US	08/829,536	28 March 1997 (28.03.97)	US	08/869,852	4 June 1997 (04.06.97)	US	08/883,036	26 June 1997 (26.06.97)	US	<b>(81) Designated States:</b> AU, CA, IL, JP, KR, MX, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
08/799,861	13 February 1997 (13.02.97)	US															
08/815,255	12 March 1997 (12.03.97)	US															
08/829,536	28 March 1997 (28.03.97)	US															
08/869,852	4 June 1997 (04.06.97)	US															
08/883,036	26 June 1997 (26.06.97)	US															
<b>(54) Title:</b> RECEPTOR THAT BINDS TRAIL  <b>(57) Abstract</b> <p>A protein designated TRAIL receptor binds the protein known as TNF-Related Apoptosis-Inducing Ligand (TRAIL). The TRAIL receptor finds use in purifying TRAIL or inhibiting activities thereof. Isolated DNA sequences encoding TRAIL-R polypeptides are provided, along with expression vectors containing the DNA sequences, and host cells transformed with such recombinant expression vectors. Antibodies that are immunoreactive with TRAIL-R are also provided.</p>																	

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## TITLE

### RECEPTOR THAT BINDS TRAIL

5

### BACKGROUND OF THE INVENTION

A protein known as TNF-related apoptosis-inducing ligand (TRAIL) is a member of the tumor necrosis factor family of ligands (Wiley et al., *Immunity*, 3:673-682, 1995). TRAIL has demonstrated the ability to induce apoptosis of certain transformed cells, including a number of different types of cancer cells as well as virally infected cells (PCT application WO 97/01633 and Wiley et al., *supra*).

Identification of receptor protein(s) that bind TRAIL would prove useful in further study of the biological activities of TRAIL. However, prior to the present invention, no receptor for TRAIL had been reported.

15

### SUMMARY OF THE INVENTION

The present invention is directed to a novel protein designated TRAIL receptor (TRAIL-R), which binds to a protein known as TNF-related apoptosis-inducing ligand (TRAIL). DNA encoding TRAIL-R, and expression vectors comprising such DNA, are provided. A method for producing TRAIL-R polypeptides comprises culturing host cells transformed with a recombinant expression vector encoding TRAIL-R, under conditions that promote expression of TRAIL-R, then recovering the expressed TRAIL-R polypeptides from the culture. Antibodies that are immunoreactive with TRAIL-R are also provided.

25

### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 presents the nucleotide sequence of a human TRAIL receptor DNA fragment, as well as the amino acid sequence encoded thereby. This DNA fragment is described in Example 3.

30

Figure 2 presents the results of the assay described in example 7. In the assay, a soluble TRAIL-R/Fc fusion protein blocked TRAIL-induced apoptosis of Jurkat cells.

Figure 3 presents the results of the experiment described in example 8. The indicated compounds were demonstrated to inhibit apoptosis of cells expressing TRAIL receptor.

35

### DETAILED DESCRIPTION OF THE INVENTION

A novel protein designated TRAIL receptor (TRAIL-R) is provided herein. TRAIL-R binds to the cytokine designated TNF-related apoptosis-inducing ligand (TRAIL). Certain uses of TRAIL-R flow from this ability to bind TRAIL, as discussed  
5 further below. TRAIL-R finds use in inhibiting biological activities of TRAIL, or in purifying TRAIL by affinity chromatography, for example.

The nucleotide sequence of the coding region of a human TRAIL receptor DNA is presented in SEQ ID NO:1. The amino acid sequence encoded by the DNA sequence of SEQ ID NO:1 is shown in SEQ ID NO:2. This sequence information identifies the TRAIL  
10 receptor protein as a member of the tumor necrosis factor receptor (TNF-R) family of receptors (reviewed in Smith et al., *Cell* 76:959-962, 1994). The extracellular domain contains cysteine rich repeats; such motifs have been reported to be important for ligand binding in other receptors of this family. TRAIL-R contains a so-called "death domain" in the cytoplasmic region; such domains in certain other receptors are associated with  
15 transduction of apoptotic signals. These and other features of the protein are discussed in more detail below.

TRAIL-R protein or immunogenic fragments thereof may be employed as immunogens to generate antibodies that are immunoreactive therewith. In one embodiment of the invention, the antibodies are monoclonal antibodies.

20 A human TRAIL-R protein was purified as described in example 1. In example 2, amino acid sequence information derived from fragments of TRAIL-R is presented. One embodiment of the invention is directed to a purified human TRAIL-R protein that is capable of binding TRAIL, wherein the TRAIL-R is characterized as comprising the amino acid sequence VPANEGD (amino acids 327 to 333 of SEQ ID NO:2). In another  
25 embodiment, the TRAIL-R additionally comprises the sequence ETLRQCFFDDFADLVPFDSWEPLMRKLGLMDNEIKVAKAEAAGHRDTLXTML (amino acids 336 to 386 of SEQ ID NO:2, with one unknown amino acid indicated as X). Also provided are TRAIL-R fragments comprising only one of these characterizing amino acid sequences.

30 The nucleotide sequence of a TRAIL-R DNA fragment, and the amino acid sequence encoded thereby, are presented in Figure 1 (SEQ ID NO:3 and SEQ ID NO:4); see example 3. The amino acid sequence presented in Figure 1 has characteristics of the so-called "death domains" found in the cytoplasmic region of certain other receptor proteins. Such domains have been reported to be associated with transduction of apoptotic  
35 signals. Cytoplasmic death domains have been identified in Fas antigen (Itoh and Nagata, *J. Biol. Chem.* 268:10932, 1993), TNF receptor type I (Tartaglia et al. *Cell* 74:845, 1993),

DR3 (Chinnaiyan et al., *Science* 274:990-992, 1996), and CAR-1 (Brojatsch et al., *Cell* 87:845-855, 1996). The role of these death domains in initiating intracellular apoptotic signaling cascades is discussed further below.

5 SEQ ID NO:1 presents the nucleotide sequence of the coding region of a human TRAIL receptor DNA, including an initiation codon (ATG) and a termination codon (TAA). The amino acid sequence encoded by the DNA of SEQ ID NO:1 is presented in SEQ ID NO:2. The fragment depicted in Figure 1 corresponds to the region of TRAIL-R that is presented as amino acids 336 to 386 in SEQ ID NO:2.

10 The TRAIL-R protein of SEQ ID NO:2 includes an N-terminal hydrophobic region that functions as a signal peptide, followed by an extracellular domain, a transmembrane region comprising amino acids 211 through 231, and a C-terminal cytoplasmic domain. Computer analysis predicts that the signal peptide corresponds to residues 1 to 51 of SEQ ID NO:2. Cleavage of the signal peptide thus would yield a mature protein comprising amino acids 52 through 440 of SEQ ID NO:2. The calculated molecular weight for a  
15 mature protein containing residues 52 to 440 of SEQ ID NO:2 is about 43 kilodaltons. The next most likely computer-predicted signal peptidase cleavage sites (in descending order) occur after amino acids 50 and 58 of SEQ ID NO:2.

In another embodiment of the invention, the N-terminal residue of a mature TRAIL-R protein is the isoleucine residue at position 56 of SEQ ID NO:2. Sequences of several  
20 tryptic digest peptide fragments of TRAIL-R were determined by a combination of N-terminal sequencing and Nano-ES MS/MS (nano electrospray tandem mass spectrometry). The N-terminal amino acid of one of the peptide fragments was the isoleucine at position 56 of SEQ ID NO:2. Since this fragment was not preceded by a trypsin cleavage site, the (Ile)56 residue may correspond to the N-terminal residue resulting from cleavage of the  
25 signal peptide.

A further embodiment of the invention is directed to mature TRAIL-R having amino acid 54 as the N-terminal residue. In one preparation of TRAIL-R (a soluble TRAIL-R/Fc fusion protein expressed in CV1-EBNA cells), the signal peptide was cleaved after residue  
30 53 of SEQ ID NO:2.

The skilled artisan will recognize that the molecular weight of particular preparations of TRAIL-R protein may differ, according to such factors as the degree of glycosylation. The glycosylation pattern of a particular preparation of TRAIL-R may vary according to the type of cells in which the protein is expressed, for example. Further, a given preparation may include multiple differentially glycosylated species of the protein.  
35 TRAIL-R polypeptides with or without associated native-pattern glycosylation are provided

herein. Expression of TRAIL-R polypeptides in bacterial expression systems, such as *E. coli*, provides non-glycosylated molecules.

5 In one embodiment, the protein is characterized by a molecular weight within the range of about 50 to 55 kilodaltons, which is the molecular weight determined for a preparation of native, full length, human TRAIL-R. Molecular weight can be determined by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

Example 1 presents one method for purifying a TRAIL-R protein. Jurkat cells are disrupted, and the subsequent purification process includes affinity chromatography (employing a chromatography matrix containing TRAIL), and reversed phase HPLC.

10 TRAIL-R polypeptides of the present invention may be purified by any suitable alternative procedure, using known protein purification techniques. In one alternative procedure, the chromatography matrix instead comprises an antibody that binds TRAIL-R. Other cell types expressing TRAIL-R (e.g., the PS-1 cells described in example 2) can be substituted for the Jurkat cells. The cells can be disrupted by any of the numerous known techniques, including freeze-thaw cycling, sonication, mechanical disruption, or by use of  
15 cell lysing agents.

The desired degree of purity depends on the intended use of the protein. A relatively high degree of purity is desired when the protein is to be administered *in vivo*, for example. Advantageously, TRAIL-R polypeptides are purified such that no protein bands  
20 corresponding to other (non-TRAIL-R) proteins are detectable upon analysis by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). It will be recognized by one skilled in the pertinent field that multiple bands corresponding to TRAIL-R protein may be visualized by SDS-PAGE, due to differential glycosylation, differential post-translational processing, and the like. TRAIL-R most preferably is purified to substantial homogeneity, as indicated  
25 by a single protein band upon analysis by SDS-PAGE. The protein band may be visualized by silver staining, Coomassie blue staining, or (if the protein is radiolabeled) by autoradiography.

The present invention encompasses TRAIL-R in various forms, including those that are naturally occurring or produced through various techniques such as procedures  
30 involving recombinant DNA technology. Such forms of TRAIL-R include, but are not limited to, fragments, derivatives, variants, and oligomers of TRAIL-R, as well as fusion proteins containing TRAIL-R or fragments thereof.

TRAIL-R may be modified to create derivatives thereof by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids,  
35 phosphate, acetyl groups and the like. Covalent derivatives of TRAIL-R may be prepared by linking the chemical moieties to functional groups on TRAIL-R amino acid side chains

or at the N-terminus or C-terminus of a TRAIL-R polypeptide. Conjugates comprising diagnostic (detectable) or therapeutic agents attached to TRAIL-R are contemplated herein, as discussed in more detail below.

5 Other derivatives of TRAIL-R within the scope of this invention include covalent or aggregative conjugates of TRAIL-R polypeptides with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. Examples of fusion proteins are discussed below in connection with TRAIL-R oligomers. Further, TRAIL-R-containing fusion proteins can comprise peptides added to facilitate purification and identification of TRAIL-R. Such peptides include, for example, poly-His or the  
10 antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *Bio/Technology* 6:1204, 1988. One such peptide is the Flag<sup>®</sup> peptide, Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys, which is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. A murine hybridoma designated 4E11 produces a  
15 monoclonal antibody that binds the Flag<sup>®</sup> peptide in the presence of certain divalent metal cations, as described in U.S. Patent 5,011,912, hereby incorporated by reference. The 4E11 hybridoma cell line has been deposited with the American Type Culture Collection under accession no. HB 9259. Monoclonal antibodies that bind the Flag<sup>®</sup> peptide are available from Eastman Kodak Co., Scientific Imaging Systems Division, New Haven,  
20 Connecticut.

Both cell membrane-bound and soluble (secreted) forms of TRAIL-R are provided herein. Soluble TRAIL-R may be identified (and distinguished from non-soluble membrane-bound counterparts) by separating intact cells expressing a TRAIL-R polypeptide from the culture medium, e.g., by centrifugation, and assaying the medium  
25 (supernatant) for the presence of the desired protein. The presence of TRAIL-R in the medium indicates that the protein was secreted from the cells and thus is a soluble form of the desired protein.

Soluble forms of receptor proteins typically lack the transmembrane region that would cause retention of the protein on the cell surface. In one embodiment of the  
30 invention, a soluble TRAIL-R polypeptide comprises the extracellular domain of the protein. A soluble TRAIL-R polypeptide may include the cytoplasmic domain, or a portion thereof, as long as the polypeptide is secreted from the cell in which it is produced. One example of a soluble TRAIL-R is a soluble human TRAIL-R comprising amino acids 52 to 210 of SEQ ID NO:2. Other soluble TRAIL-R polypeptides include, but are not limited to,  
35 polypeptides comprising amino acids x to 210 of SEQ ID NO:2, wherein x is an integer from 51 through 59.

Soluble forms of TRAIL-R possess certain advantages over the membrane-bound form of the protein. Purification of the protein from recombinant host cells is facilitated, since the soluble proteins are secreted from the cells. Further, soluble proteins are generally more suitable for certain applications, e.g., for intravenous administration.

5 TRAIL-R fragments are provided herein. Such fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative involves generating TRAIL-R fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed as the 5' and 3' primers in the PCR.

10 Examples of fragments are those comprising at least 20, or at least 30, contiguous amino acids of the sequence of SEQ ID NO:2. Fragments derived from the cytoplasmic domain find use in studies of TRAIL-R-mediated signal transduction, and in regulating cellular processes associated with transduction of biological signals. TRAIL-R polypeptide fragments also may be employed as immunogens, in generating antibodies. Particular embodiments are directed to TRAIL-R polypeptide fragments that retain the ability to bind TRAIL. Such a fragment may be a soluble TRAIL-R polypeptide, as described above.

20 Naturally occurring variants of the TRAIL-R protein of SEQ ID NO:2 are provided herein. Such variants include, for example, proteins that result from alternate mRNA splicing events or from proteolytic cleavage of the TRAIL-R protein. Alternate splicing of mRNA may, for example, yield a truncated but biologically active TRAIL-R protein, such as a naturally occurring soluble form of the protein. Variations attributable to proteolysis include, for example, differences in the N- or C-termini upon expression in different types of host cells, due to proteolytic removal of one or more terminal amino acids from the TRAIL-R protein (generally from 1-5 terminal amino acids). TRAIL-R proteins in which differences in amino acid sequence are attributable to genetic polymorphism (allelic variation among individuals producing the protein) are also contemplated herein.

30 The skilled artisan will also recognize that the position(s) at which the signal peptide is cleaved may differ from that predicted by computer program, and may vary according to such factors as the type of host cells employed in expressing a recombinant TRAIL-R polypeptide. A protein preparation may include a mixture of protein molecules having different N-terminal amino acids, resulting from cleavage of the signal peptide at more than one site. As discussed above, particular embodiments of mature TRAIL-R proteins

35



provided herein include, but are not limited to, proteins having the residue at position 51, 52, 54, 56, or 59 of SEQ ID NO:2 as the N-terminal amino acid.

Regarding the discussion herein of various domains of TRAIL-R protein, the skilled artisan will recognize that the above-described boundaries of such regions of the protein are approximate. To illustrate, the boundaries of the transmembrane region (which may be predicted by using computer programs available for that purpose) may differ from those described above. Thus, soluble TRAIL-R polypeptides in which the C-terminus of the extracellular domain differs from the residue so identified above are contemplated herein.

Other naturally occurring TRAIL-R DNAs and polypeptides include those derived from non-human species. Homologs of the human TRAIL-R of SEQ ID NO:2, from other mammalian species, are contemplated herein, for example. Probes based on the human DNA sequence of SEQ ID NO:3 or SEQ ID NO:1 may be used to screen cDNA libraries derived from other mammalian species, using conventional cross-species hybridization techniques.

TRAIL-R DNA sequences may vary from the native sequences disclosed herein. Due to the known degeneracy of the genetic code, wherein more than one codon can encode the same amino acid, a DNA sequence can vary from that shown in SEQ ID NO:1 and still encode a TRAIL-R protein having the amino acid sequence of SEQ ID NO:2. Such variant DNA sequences may result from silent mutations (e.g., occurring during PCR amplification), or may be the product of deliberate mutagenesis of a native sequence. Thus, among the DNA sequences provided herein are native TRAIL-R sequences (e.g., cDNA comprising the nucleotide sequence presented in SEQ ID NO:1) and DNA that is degenerate as a result of the genetic code to a native TRAIL-R DNA sequence.

Among the TRAIL-R polypeptides provided herein are variants of native TRAIL-R polypeptides that retain a biological activity of a native TRAIL-R. Such variants include polypeptides that are substantially homologous to native TRAIL-R, but which have an amino acid sequence different from that of a native TRAIL-R because of one or more deletions, insertions or substitutions. Particular embodiments include, but are not limited to, TRAIL-R polypeptides that comprise from one to ten deletions, insertions or substitutions of amino acid residues, when compared to a native TRAIL-R sequence. The TRAIL-R-encoding DNAs of the present invention include variants that differ from a native TRAIL-R DNA sequence because of one or more deletions, insertions or substitutions, but that encode a biologically active TRAIL-R polypeptide. One biological activity of TRAIL-R is the ability to bind TRAIL.

Nucleic acid molecules capable of hybridizing to the DNA of SEQ ID NO:1 or SEQ ID NO:3 under moderately stringent or highly stringent conditions, and which encode a biologically active TRAIL-R, are provided herein. Such hybridizing nucleic acids include, but are not limited to, variant DNA sequences and DNA derived from non-human species.  
5 e.g., non-human mammals.

Moderately stringent conditions include conditions described in, for example, Sambrook et al, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Vol. 1, pp 1.101-104. Cold Spring Harbor Laboratory Press, 1989. Conditions of moderate stringency, as defined by Sambrook et al., include use of a prewashing solution of 5X SSC, 0.5% SDS,  
10 1.0 mM EDTA (pH 8.0) and hybridization conditions of about 55°C, 5 X SSC, overnight. Highly stringent conditions include higher temperatures of hybridization and washing. One embodiment of the invention is directed to DNA sequences that will hybridize to the DNA of SEQ ID NOS:1 or 3 under highly stringent conditions, wherein said conditions include hybridization at 68°C followed by washing in 0.1X SSC/0.1% SDS at 63-68°C.

15 Certain DNAs and polypeptides provided herein comprise nucleotide or amino acid sequences, respectively, that are at least 80% identical to a native TRAIL-R sequence. Also contemplated are embodiments in which a TRAIL-R DNA or polypeptide comprises a sequence that is at least 90% identical, at least 95% identical, or at least 98% identical to a native TRAIL-R sequence. The percent identity may be determined, for example, by  
20 comparing sequence information using the GAP computer program, version 6.0 described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov  
25 and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

In particular embodiments of the invention, a variant TRAIL-R polypeptide differs  
30 in amino acid sequence from a native TRAIL-R, but is substantially equivalent to a native TRAIL-R in a biological activity. One example is a variant TRAIL-R that binds TRAIL with essentially the same binding affinity as does a native TRAIL-R. Binding affinity can be measured by conventional procedures, e.g., as described in U.S. Patent no. 5,512,457.

35 Variant amino acid sequences may comprise conservative substitution(s), meaning that one or more amino acid residues of a native TRAIL-R is replaced by a different

residue, but that the conservatively substituted TRAIL-R polypeptide retains a desired biological activity of the native protein (e.g., the ability to bind TRAIL). A given amino acid may be replaced by a residue having similar physiochemical characteristics. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other conservative substitutions, e.g., involving substitutions of entire regions having similar hydrophobicity characteristics, are well known.

In another example of variants, sequences encoding Cys residues that are not essential for biological activity can be altered to cause the Cys residues to be deleted or replaced with other amino acids, preventing formation of incorrect intramolecular disulfide bridges upon renaturation. Certain receptors of the TNF-R family contain cysteine-rich repeat motifs in their extracellular domains (Marsters et al., J. Biol. Chem. 267:5747-5750, 1992). These repeats are believed to be important for ligand binding. To illustrate, Marsters et al., *supra*, reported that soluble TNF-R type 1 polypeptides lacking one of the repeats exhibited a ten fold reduction in binding affinity for TNF $\alpha$  and TNF $\beta$ ; deletion of the second repeat resulted in a complete loss of detectable binding of the ligands. The human TRAIL-R of SEQ ID NO:2 contains two such cysteine rich repeats, the first including residues 94 through 137, and the second including residues 138 through 178. Cysteine residues within these cysteine rich domains advantageously remain unaltered in TRAIL-R variants, when retention of TRAIL-binding activity is desired.

Other variants are prepared by modification of adjacent dibasic amino acid residues, to enhance expression in yeast systems in which KEX2 protease activity is present. EP 212,914 discloses the use of site-specific mutagenesis to inactivate KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Mature human TRAIL-R contains such adjacent basic residue pairs at amino acids 72-73, 154-155, 322-323, 323-324, and 359-360 of SEQ ID NO:2. Lys-Lys pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites.

TRAIL-R polypeptides, including variants and fragments thereof, can be tested for biological activity in any suitable assay. The ability of a TRAIL-R polypeptide to bind TRAIL can be confirmed in conventional binding assays, examples of which are described below.

### Expression Systems

The present invention also provides recombinant cloning and expression vectors containing TRAIL-R DNA, as well as host cell containing the recombinant vectors. Expression vectors comprising TRAIL-R DNA may be used to prepare TRAIL-R polypeptides encoded by the DNA. A method for producing TRAIL-R polypeptides comprises culturing host cells transformed with a recombinant expression vector encoding TRAIL-R, under conditions that promote expression of TRAIL-R, then recovering the expressed TRAIL-R polypeptides from the culture. The skilled artisan will recognize that the procedure for purifying the expressed TRAIL-R will vary according to such factors as the type of host cells employed, and whether the TRAIL-R is membrane-bound or a soluble form that is secreted from the host cell.

Any suitable expression system may be employed. The vectors include a DNA encoding a TRAIL-R polypeptide, operably linked to suitable transcriptional or translational regulatory nucleotide sequences, such as those derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, an mRNA ribosomal binding site, and appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the TRAIL-R DNA sequence. Thus, a promoter nucleotide sequence is operably linked to an TRAIL-R DNA sequence if the promoter nucleotide sequence controls the transcription of the TRAIL-R DNA sequence. An origin of replication that confers the ability to replicate in the desired host cells, and a selection gene by which transformants are identified, are generally incorporated into the expression vector.

In addition, a sequence encoding an appropriate signal peptide (native or heterologous) can be incorporated into expression vectors. A DNA sequence for a signal peptide (secretory leader) may be fused in frame to the TRAIL-R sequence so that the TRAIL-R is initially translated as a fusion protein comprising the signal peptide. A signal peptide that is functional in the intended host cells promotes extracellular secretion of the TRAIL-R polypeptide. The signal peptide is cleaved from the TRAIL-R polypeptide upon secretion of TRAIL-R from the cell.

Suitable host cells for expression of TRAIL-R polypeptides include prokaryotes, yeast or higher eukaryotic cells. Mammalian or insect cells are generally preferred for use as host cells. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985). Cell-free translation

systems could also be employed to produce TRAIL-R polypeptides using RNAs derived from DNA constructs disclosed herein.

Prokaryotes include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*,  
5 *Bacillus subtilis*, *Salmonella typhimurium*, and various other species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a prokaryotic host cell, such as *E. coli*, a TRAIL-R polypeptide may include an N-terminal methionine residue to facilitate expression of the recombinant polypeptide in the prokaryotic host cell. The N-terminal Met may be cleaved from the expressed recombinant TRAIL-R polypeptide.

10 Expression vectors for use in prokaryotic host cells generally comprise one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322  
15 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. An appropriate promoter and a TRAIL-R DNA sequence are inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA).

20 Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include  $\beta$ -lactamase (penicillinase), lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EP-A-36776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor  
25 Laboratory, p. 412, 1982). A particularly useful prokaryotic host cell-expression system employs a phage  $\lambda$  P<sub>L</sub> promoter and a cI857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the  $\lambda$  P<sub>L</sub> promoter include plasmid pHUB2 (resident in *E. coli* strain JMB9, ATCC 37092) and pPLc28 (resident in *E. coli* RR1, ATCC 53082).

30 TRAIL-R alternatively may be expressed in yeast host cells, preferably from the *Saccharomyces* genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia* or *Kluyveromyces*, may also be employed. Yeast vectors will often contain an origin of replication sequence from a 2 $\mu$  yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription  
35 termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase

(Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phospho-glucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA-73,657. Another alternative is the glucose-repressible ADH2 promoter described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). Shuttle vectors replicable in both yeast and *E. coli* may be constructed by inserting DNA sequences from pBR322 for selection and replication in *E. coli* (Amp<sup>r</sup> gene and origin of replication) into the above-described yeast vectors.

The yeast  $\alpha$ -factor leader sequence may be employed to direct secretion of the TRAIL polypeptide. The  $\alpha$ -factor leader sequence is often inserted between the promoter sequence and the structural gene sequence. See, e.g., Kurjan et al., *Cell* 30:933, 1982 and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. Other leader sequences suitable for facilitating secretion of recombinant polypeptides from yeast hosts are known to those of skill in the art. A leader sequence may be modified near its 3' end to contain one or more restriction sites. This will facilitate fusion of the leader sequence to the structural gene.

Yeast transformation protocols are known to those of skill in the art. One such protocol is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978. The Hinnen et al. protocol selects for Trp<sup>+</sup> transformants in a selective medium, wherein the selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10  $\mu$ g/ml adenine and 20  $\mu$ g/ml uracil.

Yeast host cells transformed by vectors containing an ADH2 promoter sequence may be grown for inducing expression in a "rich" medium. An example of a rich medium is one consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80  $\mu$ g/ml adenine and 80  $\mu$ g/ml uracil. Derepression of the ADH2 promoter occurs when glucose is exhausted from the medium.

Mammalian or insect host cell culture systems also may be employed to express recombinant TRAIL-R polypeptides. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988). Established cell lines of mammalian origin also may be employed. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, and BHK (ATCC CRL 10)

cell lines, and the CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10: 2821, 1991).

Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* I site located in the SV40 viral origin of replication site is included.

Expression vectors for use in mammalian host cells can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983), for example. A useful system for stable high level expression of mammalian cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984 has been deposited as ATCC 39890. Additional mammalian expression vectors are described in EP-A-0367566, and in WO 91/18982. As one alternative, the vector may be derived from a retrovirus. Overexpression of full length TRAIL-R has resulted in membrane blebbing and nuclear condensation of transfected CV-1/EBNA cells, indicating that the mechanism of cell death was apoptosis. For host cells in which such TRAIL-R-mediated apoptosis occurs, a suitable apoptosis inhibitor may be included in the expression system.

To inhibit TRAIL-R-induced apoptosis of host cells expressing recombinant TRAIL-R, the cells may be co-transfected with an expression vector encoding a polypeptide that functions as an apoptosis inhibitor. Expression vectors encoding such polypeptides can be prepared by conventional procedures. Another approach involves adding an apoptosis inhibitor to the culture medium. The use of poxvirus CrmA, baculovirus P35, a C-terminal fragment of FADD, and the tripeptide derivative zVAD-fmk, to reduce death of host cells is illustrated in examples 6 and 8.

zVAD-fmk (benzyloxycarbonyl-Val-Ala-Asp-fluoromethylketone) is a tripeptide based compound, available from Enzyme System Products, Dublin, California. As

illustrated in example 8, zVAD-fmk may be added to the medium in which cells expressing TRAIL-R are cultured.

5 The 38-kilodalton cowpox-derived protein that was subsequently designated CrmA is described in Pickup et al. (*Proc. Natl. Acad. Sci. USA* 83:7698-7702, 1986; hereby incorporated by reference). Sequence information for CrmA is presented in Figure 4 of Pickup et al., *supra*. One approach to producing and purifying CrmA protein is described in Ray et al. (*Cell*, 69:597-604, 1992; hereby incorporated by reference).

10 A 35-kilodalton protein encoded by *Autographa californica* nuclear polyhedrosis virus, a baculovirus, is described in Friesen and Miller (*J. Virol.* 61:2264-2272, 1987; hereby incorporated by reference). Sequence information for this protein, designated baculovirus p35 herein, is presented in Figure 5 of Friesen and Miller, *supra*.

15 The death domain-containing cytoplasmic protein FADD (also known as MORT1) is described in Boldin et al. (*J. Biol. Chem.* 270:7795-7798, 1995; hereby incorporated by reference). FADD has been reported to associate, directly or indirectly, with the cytoplasmic death domain of certain receptors that mediate apoptosis (Boldin et al., *Cell* 85:803-815, June 1996; Hsu et al., *Cell* 84:299-308, January 1996).

20 In one embodiment of the present invention, truncated FADD polypeptides that include the death domain (located in the C-terminal portion of the protein), but lack the N-terminal region to which apoptosis effector functions have been attributed, are employed to reduce apoptosis. The use of certain FADD deletion mutant polypeptides, truncated at the N-terminus, to inhibit death of cells expressing other apoptosis-inducing receptors, is described in Hsu et al. (*Cell* 84:299-308, 1996; hereby incorporated by reference).

25 This approach is illustrated in example 8, which employs one suitable FADD-dominant negative (FADD-DN) polypeptide, having an amino acid sequence corresponding to amino acids 117 through 245 of the MORT1 amino acid sequence presented in Boldin et al. (*J. Biol. Chem.* 270:7795-7798, 1995). In example 8, cells were co-transfected with a TRAIL-R-encoding expression vector, and with an expression vector encoding the above-described Flag® peptide, fused to the N-terminus of the FADD-DN polypeptide.

30 While not wishing to be bound by theory, one possible explanation is that the C-terminal fragments of FADD associate with the intracellular death domain of the receptor, but lack the N-terminal portion of the protein that is necessary for effecting apoptosis (Hsu et al., *Cell* 84:299-308, January 1996; Boldin et al., *Cell* 85:803-815, June 1996). The truncated FADD thereby may block association of endogenous, full length FADD with the receptor's death domain; consequently, the apoptosis that would be initiated by such  
35 endogenous FADD is inhibited.



Other apoptosis inhibitors useful in expression systems of the present invention can be identified in conventional assay procedures. One such assay, in which compounds are tested for the ability to reduce apoptosis of cells expressing TRAIL-R, is described in example 8.

5 Poxvirus CrmA, baculovirus P35, and zVAD-fmk are viral caspase inhibitors. Other caspase inhibitors may be tested for the ability to reduce TRAIL-R-mediated cell death.

The use of CrmA, baculovirus p35, and certain peptide derivatives (including zVAD-fmk) as inhibitors of apoptosis in particular cells/systems is discussed in Sarin et al.  
10 (*J. Exp. Med.* 184:2445-2450, Dec.-1996; hereby incorporated by reference). The role of interleukin-1 $\beta$  converting enzyme (ICE) family proteases in signal transduction cascades leading to programmed cell death, and the use of inhibitors of such proteases to block apoptosis, is discussed in Sarin et al., *supra*, and Muzio et al., *Cell* 85:817-827, 1996).

Apoptosis inhibitors generally need not be employed for expression of TRAIL-R  
15 polypeptides lacking the cytoplasmic domain (i.e., lacking the region of the protein involved in signal transduction). Thus, expression systems for producing soluble TRAIL-R polypeptides comprising only the extracellular domain (or a fragment thereof) need not include one of the above-described apoptosis inhibitors.

Regarding signal peptides that may be employed in producing TRAIL-R, the native  
20 signal peptide of TRAIL-R may be replaced by a heterologous signal peptide or leader sequence, if desired. The choice of signal peptide or leader may depend on factors such as the type of host cells in which the recombinant TRAIL-R is to be produced. To illustrate, examples of heterologous signal peptides that are functional in mammalian host cells include the signal sequence for interleukin-7 (IL-7) described in United States Patent  
25 4,965,195, the signal sequence for interleukin-2 receptor described in Cosman et al., *Nature* 312:768 (1984); the interleukin-4 receptor signal peptide described in EP 367,566; the type I interleukin-1 receptor signal peptide described in U.S. Patent 4,968,607; and the type II interleukin-1 receptor signal peptide described in EP 460,846.

### 30 Oligomeric Forms of TRAIL-R

Encompassed by the present invention are oligomers that contain TRAIL-R polypeptides. TRAIL-R oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers.

One embodiment of the invention is directed to oligomers comprising multiple  
35 TRAIL-R polypeptides joined *via* covalent or non-covalent interactions between peptide moieties fused to the TRAIL-R polypeptides. Such peptides may be peptide linkers

(spacers), or peptides that have the property of promoting oligomerization. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of TRAIL-R polypeptides attached thereto, as described in more detail below.

5 In particular embodiments, the oligomers comprise from two to four TRAIL-R polypeptides. The TRAIL-R moieties of the oligomer may be soluble polypeptides, as described above.

As one alternative, a TRAIL-R oligomer is prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al. (*PNAS USA* 88:10535, 1991); Byrn et al. (*Nature* 344:677, 1990); and Hollenbaugh and Aruffo ("Construction of Immunoglobulin Fusion Proteins", in *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1 - 10.19.11, 1992).

15 One embodiment of the present invention is directed to a TRAIL-R dimer comprising two fusion proteins created by fusing TRAIL-R to the Fc region of an antibody. A gene fusion encoding the TRAIL-R/Fc fusion protein is inserted into an appropriate expression vector. TRAIL-R/Fc fusion proteins are expressed in host cells transformed with the recombinant expression vector, and allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield divalent TRAIL-R.

Provided herein are fusion proteins comprising a TRAIL-R polypeptide fused to an Fc polypeptide derived from an antibody. DNA encoding such fusion proteins, as well as dimers containing two fusion proteins joined *via* disulfide bonds between the Fc moieties thereof, are also provided. The term "Fc polypeptide" as used herein includes native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization are also included.

One suitable Fc polypeptide, described in PCT application WO 93/10151 (hereby incorporated by reference), is a single chain polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Patent 5,457,035 and in Baum et al., (*EMBO J.* 13:3992-4001, 1994). The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and

amino acid 22 has been changed from Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

5 In other embodiments, TRAIL-R may be substituted for the variable portion of an antibody heavy or light chain. If fusion proteins are made with both heavy and light chains of an antibody, it is possible to form a TRAIL-R oligomer with as many as four TRAIL-R extracellular regions.

10 Alternatively, the oligomer is a fusion protein comprising multiple TRAIL-R polypeptides, with or without peptide linkers (spacer peptides). Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233, which are hereby incorporated by reference. A DNA sequence encoding a desired peptide linker may be inserted between, and in the same reading frame as, the DNA sequences encoding TRAIL-R, using any suitable conventional technique. For example, a chemically synthesized oligonucleotide encoding the linker may be ligated between sequences encoding TRAIL-R. In particular embodiments, a fusion protein comprises from two to 15 four soluble TRAIL-R polypeptides, separated by peptide linkers.

Another method for preparing oligomeric TRAIL-R involves use of a leucine zipper. Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., *Science* 240:1759, 1988), and have since been found in a 20 variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe et al. (*FEBS Letters* 344:191, 1994), hereby incorporated by 25 reference. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow et al. (*Semin. Immunol.* 6:267-278, 1994). Recombinant fusion proteins comprising a soluble TRAIL-R polypeptide fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric TRAIL-R that forms is recovered from the culture supernatant.

30 Oligomeric TRAIL-R has the property of bivalent, trivalent, etc. binding sites for TRAIL. The above-described fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns. DNA sequences encoding oligomeric TRAIL-R, or encoding fusion proteins useful in preparing TRAIL-R oligomers, are provided herein.

35

### Assays

TRAIL-R proteins (including fragments, variants, oligomers, and other forms of TRAIL-R) may be tested for the ability to bind TRAIL in any suitable assay, such as a conventional binding assay. To illustrate, TRAIL-R may be labeled with a detectable reagent (e.g., a radionuclide, chromophore, enzyme that catalyzes a colorimetric or fluorometric reaction, and the like). The labeled TRAIL-R is contacted with cells expressing TRAIL. The cells then are washed to remove unbound labeled TRAIL-R, and the presence of cell-bound label is determined by a suitable technique, chosen according to the nature of the label.

One example of a binding assay procedure is as follows. A recombinant expression vector containing TRAIL cDNA is constructed, e.g., as described in PCT application WO 97/01633, hereby incorporated by reference. DNA and amino acid sequence information for human and mouse TRAIL is presented in WO 97/01633. TRAIL comprises an N-terminal cytoplasmic domain, a transmembrane region, and a C-terminal extracellular domain. CV1-EBNA-1 cells in 10 cm<sup>2</sup> dishes are transfected with the recombinant expression vector. CV1-EBNA-1 cells (ATCC CRL 10478) constitutively express EBV nuclear antigen-1 driven from the CMV immediate-early enhancer/promoter. CV1-EBNA-1 was derived from the African Green Monkey kidney cell line CV-1 (ATCC CCL 70), as described by McMahan et al. (*EMBO J.* 10:2821, 1991).

The transfected cells are cultured for 24 hours, and the cells in each dish then are split into a 24-well plate. After culturing an additional 48 hours, the transfected cells (about 4 x 10<sup>4</sup> cells/well) are washed with BM-NFDM, which is binding medium (RPMI 1640 containing 25 mg/ml bovine serum albumin, 2 mg/ml sodium azide, 20 mM Hepes pH 7.2) to which 50 mg/ml nonfat dry milk has been added. The cells then are incubated for 1 hour at 37°C with various concentrations of a soluble TRAIL-R/Fc fusion protein. Cells then are washed and incubated with a constant saturating concentration of a <sup>125</sup>I-mouse anti-human IgG in binding medium, with gentle agitation for 1 hour at 37°C. After extensive washing, cells are released *via* trypsinization.

The mouse anti-human IgG employed above is directed against the Fc region of human IgG and can be obtained from Jackson ImmunoResearch Laboratories, Inc., West Grove, PA. The antibody is radioiodinated using the standard chloramine-T method. The antibody will bind to the Fc portion of any TRAIL-R/Fc protein that has bound to the cells. In all assays, non-specific binding of <sup>125</sup>I-antibody is assayed in the absence of TRAIL-R/Fc, as well as in the presence of TRAIL-R/Fc and a 200-fold molar excess of unlabeled mouse anti-human IgG antibody.

Cell-bound  $^{125}\text{I}$ -antibody is quantified on a Packard Autogamma counter. Affinity calculations (Scatchard, *Ann. N.Y. Acad. Sci.* 51:660, 1949) are generated on RS/1 (BBN Software, Boston, MA) run on a Microvax computer.

Another type of suitable binding assay is a competitive binding assay. To illustrate, biological activity of a TRAIL-R variant may be determined by assaying for the variant's ability to compete with a native TRAIL-R for binding to TRAIL.

Competitive binding assays can be performed by conventional methodology. Reagents that may be employed in competitive binding assays include radiolabeled TRAIL-R and intact cells expressing TRAIL (endogenous or recombinant) on the cell surface. For example, a radiolabeled soluble TRAIL-R fragment can be used to compete with a soluble TRAIL-R variant for binding to cell surface TRAIL. Instead of intact cells, one could substitute a soluble TRAIL/Fc fusion protein bound to a solid phase through the interaction of Protein A or Protein G (on the solid phase) with the Fc moiety. Chromatography columns that contain Protein A and Protein G include those available from Pharmacia Biotech, Inc., Piscataway, NJ. Another type of competitive binding assay utilizes radiolabeled soluble TRAIL, such as a soluble TRAIL/Fc fusion protein, and intact cells expressing TRAIL-R. Qualitative results can be obtained by competitive autoradiographic plate binding assays, while Scatchard plots (Scatchard, *Ann. N.Y. Acad. Sci.* 51:660, 1949) may be utilized to generate quantitative results.

Another type of assay for biological activity involves testing a TRAIL-R polypeptide for the ability to block TRAIL-mediated apoptosis of target cells, such as the human leukemic T-cell line known as Jurkat cells, for example. TRAIL-mediated apoptosis of the cell line designated Jurkat clone E6-1 (ATCC TIB 152) is demonstrated in assay procedures described in PCT application WO 97/01633, hereby incorporated by reference.

#### Uses of TRAIL-R

Uses of TRAIL-R include, but are not limited to, the following. Certain of these uses of TRAIL-R flow from its ability to bind TRAIL.

TRAIL-R finds use as a protein purification reagent. TRAIL-R polypeptides may be attached to a solid support material and used to purify TRAIL proteins by affinity chromatography. In particular embodiments, a TRAIL-R polypeptide (in any form described herein that is capable of binding TRAIL) is attached to a solid support by conventional procedures. As one example, chromatography columns containing functional groups that will react with functional groups on amino acid side chains of proteins are available (Pharmacia Biotech, Inc., Piscataway, NJ). In an alternative, a TRAIL-R/Fc

protein is attached to Protein A- or Protein G-containing chromatography columns through interaction with the Fc moiety.

TRAIL-R proteins also find use in measuring the biological activity of TRAIL proteins in terms of their binding affinity for TRAIL-R. TRAIL-R proteins thus may be employed by those conducting "quality assurance" studies, e.g., to monitor shelf life and stability of TRAIL protein under different conditions. To illustrate, TRAIL-R may be employed in a binding affinity study to measure the biological activity of a TRAIL protein that has been stored at different temperatures, or produced in different cell types. TRAIL-R also may be used to determine whether biological activity is retained after modification of a TRAIL protein (e.g., chemical modification, truncation, mutation, etc.). The binding affinity of the modified TRAIL protein for TRAIL-R is compared to that of an unmodified TRAIL protein to detect any adverse impact of the modifications on biological activity of TRAIL. The biological activity of a TRAIL protein thus can be ascertained before it is used in a research study, for example.

TRAIL-R also finds use in purifying or identifying cells that express TRAIL on the cell surface. TRAIL-R polypeptides are bound to a solid phase such as a column chromatography matrix or a similar suitable substrate. For example, magnetic microspheres can be coated with TRAIL-R and held in an incubation vessel through a magnetic field. Suspensions of cell mixtures containing TRAIL-expressing cells are contacted with the solid phase having TRAIL-R thereon. Cells expressing TRAIL on the cell surface bind to the fixed TRAIL-R, and unbound cells then are washed away.

Alternatively, TRAIL-R can be conjugated to a detectable moiety, then incubated with cells to be tested for TRAIL expression. After incubation, unbound labeled TRAIL-R is removed and the presence or absence of the detectable moiety on the cells is determined.

In a further alternative, mixtures of cells suspected of containing TRAIL<sup>+</sup> cells are incubated with biotinylated TRAIL-R. Incubation periods are typically at least one hour in duration to ensure sufficient binding. The resulting mixture then is passed through a column packed with avidin-coated beads, whereby the high affinity of biotin for avidin provides binding of the desired cells to the beads. Procedures for using avidin-coated beads are known (see Berenson, et al. *J. Cell. Biochem.*, 10D:239, 1986). Washing to remove unbound material, and the release of the bound cells, are performed using conventional methods.

TRAIL-R polypeptides also find use as carriers for delivering agents attached thereto to cells bearing TRAIL. Cells expressing TRAIL include those identified in Wiley et al. (*Immunity*, 3:673-682, 1995). TRAIL-R proteins thus can be used to deliver

diagnostic or therapeutic agents to such cells (or to other cell types found to express TRAIL on the cell surface) in *in vitro* or *in vivo* procedures.

Detectable (diagnostic) and therapeutic agents that may be attached to a TRAIL-R polypeptide include, but are not limited to, toxins, other cytotoxic agents, drugs, 5 radionuclides, chromophores, enzymes that catalyze a colorimetric or fluorometric reaction, and the like, with the particular agent being chosen according to the intended application. Among the toxins are ricin, abrin, diphtheria toxin, *Pseudomonas aeruginosa* exotoxin A, ribosomal inactivating proteins, mycotoxins such as trichothecenes, and derivatives and fragments (e.g., single chains) thereof. Radionuclides suitable for diagnostic use include, 10 but are not limited to,  $^{123}\text{I}$ ,  $^{131}\text{I}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{111}\text{In}$ , and  $^{76}\text{Br}$ . Examples of radionuclides suitable for therapeutic use are  $^{131}\text{I}$ ,  $^{211}\text{At}$ ,  $^{77}\text{Br}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{212}\text{Pb}$ ,  $^{212}\text{Bi}$ ,  $^{109}\text{Pd}$ ,  $^{64}\text{Cu}$ , and  $^{67}\text{Cu}$ .

Such agents may be attached to the TRAIL-R by any suitable conventional procedure. TRAIL-R, being a protein, comprises functional groups on amino acid side 15 chains that can be reacted with functional groups on a desired agent to form covalent bonds, for example. Alternatively, the protein or agent may be derivatized to generate or attach a desired reactive functional group. The derivatization may involve attachment of one of the bifunctional coupling reagents available for attaching various molecules to proteins (Pierce Chemical Company, Rockford, Illinois). A number of techniques for 20 radiolabeling proteins are known. Radionuclide metals may be attached to TRAIL-R by using a suitable bifunctional chelating agent, for example.

Conjugates comprising TRAIL-R and a suitable diagnostic or therapeutic agent (preferably covalently linked) are thus prepared. The conjugates are administered or otherwise employed in an amount appropriate for the particular application.

25 TRAIL-R DNA and polypeptides of the present invention may be used in developing treatments for any disorder mediated (directly or indirectly) by defective, or insufficient amounts of, TRAIL-R. TRAIL-R polypeptides may be administered to a mammal afflicted with such a disorder. Alternatively, a gene therapy approach may be taken. Disclosure herein of native TRAIL-R nucleotide sequences permits the detection of 30 defective TRAIL-R genes, and the replacement thereof with normal TRAIL-R-encoding genes. Defective genes may be detected in *in vitro* diagnostic assays, and by comparison of a native TRAIL-R nucleotide sequence disclosed herein with that of a TRAIL-R gene derived from a person suspected of harboring a defect in this gene.

Another use of the protein of the present invention is as a research tool for studying 35 the biological effects that result from inhibiting TRAIL/TRAIL-R interactions on different

cell types. TRAIL-R polypeptides also may be employed in *in vitro* assays for detecting TRAIL or TRAIL-R or the interactions thereof.

TRAIL-R may be employed in inhibiting a biological activity of TRAIL, in *in vitro* or *in vivo* procedures. A purified TRAIL-R polypeptide may be used to inhibit binding of  
5 TRAIL to endogenous cell surface TRAIL-R. Biological effects that result from the binding of TRAIL to endogenous receptors thus are inhibited. Various forms of TRAIL-R may be employed, including, for example, the above-described TRAIL-R fragments, oligomers, derivatives, and variants that are capable of binding TRAIL. In one embodiment, a soluble TRAIL-R is employed to inhibit a biological activity of TRAIL,  
10 e.g., to inhibit TRAIL-mediated apoptosis of particular cells.

TRAIL-R may be administered to a mammal to treat a TRAIL-mediated disorder. Such TRAIL-mediated disorders include conditions caused (directly or indirectly) or exacerbated by TRAIL.

TRAIL-R may be useful for treating thrombotic microangiopathies. One such  
15 disorder is thrombotic thrombocytopenic purpura (TTP) (Kwaan, H.C., *Semin. Hematol.*, 24:71, 1987; Thompson et al., *Blood*, 80:1890, 1992). Increasing TTP-associated mortality rates have been reported by the U.S. Centers for Disease Control (Torok et al., *Am. J. Hematol.* 50:84, 1995).

Plasma from patients afflicted with TTP (including HIV<sup>+</sup> and HIV<sup>-</sup> patients) induces  
20 apoptosis of human endothelial cells of dermal microvascular origin, but not large vessel origin (Laurence et al., *Blood*, 87:3245, April 15, 1996). Plasma of TTP patients thus is thought to contain one or more factors that directly or indirectly induce apoptosis. As described in PCT application WO 97/01633 (hereby incorporated by reference), TRAIL is present in the serum of TTP patients, and may play a role in inducing apoptosis of  
25 microvascular endothelial cells.

Another thrombotic microangiopathy is hemolytic-uremic syndrome (HUS) (Moake, J.L., *Lancet*, 343:393, 1994; Melnyk et al., *Arch. Intern. Med.*, 155:2077, 1995; Thompson et al., *supra*). One embodiment of the invention is directed to use of  
30 TRAIL-R to treat the condition that is often referred to as "adult HUS" (even though it can strike children as well). A disorder known as childhood/diarrhea-associated HUS differs in etiology from adult HUS.

Other conditions characterized by clotting of small blood vessels may be treated using TRAIL-R. Such conditions include but are not limited to the following. Cardiac  
35 problems seen in about 5-10% of pediatric AIDS patients are believed to involve clotting of small blood vessels. Breakdown of the microvasculature in the heart has been reported in



multiple sclerosis patients. As a further example, treatment of systemic lupus erythematosus (SLE) is contemplated.

5 In one embodiment, a patient's blood or plasma is contacted with TRAIL-R *ex vivo*. The TRAIL-R may be bound to a suitable chromatography matrix by conventional procedures. The patient's blood or plasma flows through a chromatography column containing TRAIL-R bound to the matrix, before being returned to the patient. The immobilized receptor binds TRAIL, thus removing TRAIL protein from the patient's blood.

10 Alternatively, TRAIL-R may be administered *in vivo* to a patient afflicted with a thrombotic microangiopathy. In one embodiment, a soluble form of TRAIL-R is administered to the patient.

The present invention thus provides a method for treating a thrombotic microangiopathy, involving use of an effective amount of TRAIL-R. A TRAIL-R polypeptide may be employed in *in vivo* or *ex vivo* procedures, to inhibit TRAIL-mediated damage to (e.g., apoptosis of) microvascular endothelial cells.

15 TRAIL-R may be employed in conjunction with other agents useful in treating a particular disorder. In an *in vitro* study reported by Laurence et al. (*Blood* 87:3245, 1996), some reduction of TTP plasma-mediated apoptosis of microvascular endothelial cells was achieved by using an anti-Fas blocking antibody, aurintricarboxylic acid, or normal plasma depleted of cryoprecipitate.

20 Thus, a patient may be treated with an agent that inhibits Fas-ligand-mediated apoptosis of endothelial cells, in combination with an agent that inhibits TRAIL-mediated apoptosis of endothelial cells. In one embodiment, TRAIL-R and an anti-FAS blocking antibody are both administered to a patient afflicted with a disorder characterized by thrombotic microangiopathy, such as TTP or HUS. Examples of blocking monoclonal antibodies directed against Fas antigen (CD95) are described in PCT application publication number WO 95/10540, hereby incorporated by reference.

25 Another embodiment of the present invention is directed to the use of TRAIL-R to reduce TRAIL-mediated death of T cells in HIV-infected patients. The role of T cell apoptosis in the development of AIDS has been the subject of a number of studies (see, for example, Meyaard et al., *Science* 257:217-219, 1992; Groux et al., *J Exp. Med.* 175:331, 1992; and Oyaizu et al., in *Cell Activation and Apoptosis in HIV Infection*, Andrieu and Lu, Eds., Plenum Press, New York, 1995, pp. 101-114). Certain investigators have studied the role of Fas-mediated apoptosis; the involvement of interleukin-1 $\beta$ -converting enzyme (ICE) also has been explored (Estaquier et al., *Blood* 87:4959-4966, 1996; Mitra et

Examples of such cells are the Jurkat cells and PS1 cells described in example 2 below. Alternatively, blocking antibodies may be identified in assays for the ability to inhibit a biological effect that results from binding of TRAIL to target cells. Antibodies may be assayed for the ability to inhibit TRAIL-mediated lysis of Jurkat cells, for example.

5        Such an antibody may be employed in an *in vitro* procedure, or administered *in vivo* to inhibit a TRAIL-R-mediated biological activity. Disorders caused or exacerbated (directly or indirectly) by the interaction of TRAIL with cell surface TRAIL receptor thus may be treated. A therapeutic method involves *in vivo* administration of a blocking  
10        antibody to a mammal in an amount effective in inhibiting a TRAIL-mediated biological activity. Disorders caused or exacerbated by TRAIL, directly or indirectly, are thus treated. Monoclonal antibodies are generally preferred for use in such therapeutic methods. In one embodiment, an antigen-binding antibody fragment is employed.

      A blocking antibody directed against TRAIL-R may be substituted for TRAIL-R in the above-described method of treating thrombotic microangiopathy, e.g., in treating TTP  
15        or HUS. The antibody is administered *in vivo*, to inhibit TRAIL-mediated damage to (e.g., apoptosis of) microvascular endothelial cells.

      Antibodies raised against TRAIL-R may be screened for agonistic (i.e., ligand-mimicking) properties. Such antibodies, upon binding to cell surface TRAIL-R, induce biological effects (e.g., transduction of biological signals) similar to the biological effects  
20        induced when TRAIL binds to cell surface TRAIL-R. Agonistic antibodies may be used to induce apoptosis of certain cancer cells or virally infected cells, as has been reported for TRAIL. The ability of TRAIL to kill cancer cells (including but not limited to leukemia, lymphoma, and melanoma cells) and virally infected cells is described in Wiley et al. (*Immunity* 3:673-682, 1995); and in PCT application WO 97/01633.

25        Compositions comprising an antibody that is directed against TRAIL-R, and a physiologically acceptable diluent, excipient, or carrier, are provided herein. Suitable components of such compositions are as described above for compositions containing TRAIL-R proteins.

      Also provided herein are conjugates comprising a detectable (e.g., diagnostic) or  
30        therapeutic agent, attached to an antibody directed against TRAIL-R. Examples of such agents are presented above. The conjugates find use in *in vitro* or *in vivo* procedures.

#### Nucleic Acids

      The present invention provides TRAIL-R nucleic acids. Such nucleic acids include,  
35        but are not limited to, DNA encoding the peptide described in example 2. Such DNAs can be identified from knowledge of the genetic code. Other nucleic acids of the present

invention include isolated DNAs comprising the nucleotide sequence presented in SEQ ID NO:1 or SEQ ID NO:3.

The present invention provides isolated nucleic acids useful in the production of TRAIL-R polypeptides, e.g., in the recombinant expression systems discussed above. Such nucleic acids include, but are not limited to, the human TRAIL-R DNA of SEQ ID NO:1. Nucleic acid molecules of the present invention include TRAIL-R DNA in both single-stranded and double-stranded form, as well as the RNA complement thereof. TRAIL-R DNA includes, for example, cDNA, genomic DNA, chemically synthesized DNA, DNA amplified by PCR, and combinations thereof. Genomic DNA may be isolated by conventional techniques, e.g., using the cDNA of SEQ ID NO:1 or 3, or a suitable fragment thereof, as a probe.

DNAs encoding TRAIL-R in any of the forms contemplated herein (e.g., full length TRAIL-R or fragments thereof) are provided. Particular embodiments of TRAIL-R-encoding DNAs include a DNA encoding the full length human TRAIL-R of SEQ ID NO:2 (including the N-terminal signal peptide), and a DNA encoding a full length mature human TRAIL-R. Other embodiments include DNA encoding a soluble TRAIL-R (e.g., encoding the extracellular domain of the protein of SEQ ID NO:2, either with or without the signal peptide).

One embodiment of the invention is directed to fragments of TRAIL-R nucleotide sequences comprising at least about 17 contiguous nucleotides of a TRAIL-R DNA sequence. In other embodiments, a DNA fragment comprises at least 30, or at least 60, contiguous nucleotides of a TRAIL-R DNA sequence. Nucleic acids provided herein include DNA and RNA complements of said fragments, along with both single-stranded and double-stranded forms of the TRAIL-R DNA.

Among the uses of TRAIL-R nucleic acid fragments is use as probes or primers. Using knowledge of the genetic code in combination with the amino acid sequences set forth in example 2, sets of degenerate oligonucleotides can be prepared. Such oligonucleotides find use as primers, e.g., in polymerase chain reactions (PCR), whereby TRAIL-R DNA fragments are isolated and amplified.

Other useful fragments of the TRAIL-R nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target TRAIL-R mRNA (sense) or TRAIL-R DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of TRAIL-R DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to about 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence

encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of TRAIL-R proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10448, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO<sub>4</sub>-mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its

corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

The following examples are provided to further illustrate particular embodiments of the invention, and are not to be construed as limiting the scope of the present invention.

#### 10                    **EXAMPLE 1: Purification of TRAIL-R Protein**

A human TRAIL receptor (TRAIL-R) protein was prepared by the following procedure. Trail-R was isolated from the cell membranes of Jurkat cells, a human acute T leukemia cell line. Jurkat cells were chosen because a specific band can be affinity precipitated from surface-biotinylated Jurkat cells, using Flag®-TRAIL covalently coupled to affi-gel beads (Biorad Laboratories, Richmond, CA). The precipitated band has a molecular weight of about 52 kD. A minor specific band of about 42 kD also was present, possibly accounting for a proteolytic breakdown product or a less glycosylated form of TRAIL-R.

Approximately 50 billion Jurkat cells were harvested by centrifugation (80 ml of cell pellet), washed once with PBS, then shock frozen on liquid nitrogen. Plasma membranes were isolated according to method number three described in Maeda et al., *Biochim. et Biophys. Acta*, 731:115, 1983; hereby incorporated by reference) with five modifications:

1. The following protease inhibitors were included in all solutions at the indicated concentrations: Aprotinin, 150 nM; EDTA, 5 mM; Leupeptin, 1 μM; pA-PMSF, 20 μM; Pefabloc, 500 μM; Pepstatin A, 1 μM; PMSF, 500 μM.
2. Dithiothreitol was not used.
3. DNAase was not used in the homogenization solution.
4. 1.25 ml of homogenization buffer was used per ml of cell pellet.
5. The homogenization was accomplished by five passages through a ground glass dounce homogenizer.

After isolation of the cell membranes, proteins were solubilized by resuspending the isolated membranes in 50 ml PBS containing 1% octylglucoside and all of the above mentioned protease inhibitors at the above indicated concentrations. The resulting solution was then repeatedly vortexed during a thirty-minute incubation at 4°C. The solution was then centrifuged at 20,000 rpm in an SW28 rotor in an LE-80 Beckman ultracentrifuge

(Beckman Instruments, Inc., Palo Alto, CA) at 4°C for 30 minutes to obtain the supernatant (the membrane extract).

### Chromatography

5        The first step of purification of TRAIL-R out of the membrane extract prepared above was affinity chromatography. The membrane extract was first applied to an anti-Flag® M2 affi-gel column (10 mg of monoclonal antibody M2 coupled to 2 ml of Affi-gel beads) to adsorb any nonspecifically binding material. The flow-through was then applied to a Flag®-TRAIL affi-gel column (10 mg of recombinant protein coupled to 1 ml of affi-gel beads).

10        The Affi-gel support is an N-hydroxysuccinimide ester of a derivatized, crosslinked agarose gel bead (available from Biorad Laboratories, Richmond, CA). As discussed above, the Flag® peptide, Asp-Tyr-Lys-Asp-Asp-Asp-Lys, provides an epitope reversibly bound by specific monoclonal antibodies, enabling rapid assay and facile  
15        purification of expressed recombinant protein. M2 is a monoclonal antibody that binds Flag®. Monoclonal antibodies that bind the Flag® peptide, as well as other reagents for preparing and using Flag® fusion proteins, are available from Eastman Kodak Co., Scientific Imaging Systems Division, New Haven, Connecticut. Preparation of Flag®-TRAIL fusion proteins (comprising Flag® fused to a soluble TRAIL polypeptide) is further  
20        described in PCT application WO 97/01633, hereby incorporated by reference.

The column was washed with 25 ml of each of the following buffers, in the order indicated:

1. PBS containing 1% octylglucoside
2. PBS
- 25        3. PBS containing an additional 200 mM NaCl
4. PBS

The bound material was eluted with 50 mM Na Citrate (pH 3) in 1 ml fractions and immediately neutralized with 300 µl of 1 M Tris-HCl (pH 8.5) per fraction. The TRAIL-binding activity of each fraction was determined by a TRAIL-R-specific ELISA as  
30        described below. Fractions with high TRAIL-binding activity were pooled, brought to 0.1 % Trifluoroacetic acid (TFA), and subsequently chromatographed on a capillary reversed-phase HPLC column [500 µm internal diameter X 25 cm fused silicone capillary column packed with 5 µm Vydac C<sub>4</sub> material (Vydac, Hesperia, CA)] using a linear gradient (2% per minute) from 0% to 100% acetonitrile in water containing 0.1% TFA. Fractions  
35        containing high TRAIL-binding activity are then determined as above, pooled, and, if desired, lyophilized.

**TRAIL-R-specific ELISA:**

Serial dilutions of TRAIL-R-containing samples (in 50 mM NaHCO<sub>3</sub>, brought to pH 9 with NaOH) were coated onto Linbro/Titertek 96 well flat bottom E.I.A. microtitration plates (ICN Biomedicals Inc., Aurora, OH) at 100 µl/well. After incubation at 4°C for 16 hours, the wells were washed six times with 200 µl PBS containing 0.05% Tween-20 (PBS-Tween). The wells were then incubated with Flag®-TRAIL at 1 µg/ml in PBS-Tween with 5% fetal calf serum (FCS) for 90 minutes (100 µl per well), followed by washing as above. Next, each well was incubated with the anti-Flag® monoclonal antibody M2 at 1 µg/ml in PBS-Tween containing 5% FCS for 90 minutes (100 µl per well), followed by washing as above. Subsequently, wells were incubated with a polyclonal goat anti-mIgG1-specific horseradish peroxidase-conjugated antibody (a 1:5000 dilution of the commercial stock in PBS-Tween containing 5% FCS) for 90 minutes (100 µl per well). The HRP-conjugated antibody was obtained from Southern Biotechnology Associates, Inc., Birmingham, Alabama. Wells then were washed six times, as above.

For development of the ELISA, a substrate mix [100 µl per well of a 1:1 premix of the TMB Peroxidase Substrate and Peroxidase Solution B (Kirkegaard Perry Laboratories, Gaithersburg, Maryland)] was added to the wells. After sufficient color reaction, the enzymatic reaction was terminated by addition of 2 N H<sub>2</sub>SO<sub>4</sub> (50 µl per well). Color intensity (indicating TRAIL-binding activity) was determined by measuring extinction at 450 nm on a V Max plate reader (Molecular Devices, Sunnyvale, CA).

**EXAMPLE 2: Amino Acid Sequence**

(a) TRAIL-R purified from Jurkat cells

TRAIL-R protein isolated from Jurkat cells was digested with trypsin, using conventional procedures. Amino acid sequence analysis was conducted on one of the peptide fragments produced by the tryptic digest. The fragment was found to contain the following sequence, which corresponds to amino acids 327 to 333 of the sequence presented in SEQ ID NO:2: VPANEGD.

(b) TRAIL-R purified from PS-1 cells

TRAIL-R protein was also isolated from PS-1 cells. PS-1 is a human B cell line that spontaneously arose after lethal irradiation of human peripheral blood lymphocytes (PBLs). The TRAIL-R protein was digested with trypsin, using conventional procedures. Amino acid sequence analysis was conducted on peptide fragments that resulted from the tryptic digest. One of the fragments was found to contain the following sequence, which,

subcutaneously or intraperitoneally. Ten to twelve days later, the immunized animals are boosted with additional TRAIL-R emulsified in incomplete Freund's adjuvant. Mice are periodically boosted thereafter on a weekly to bi-weekly immunization schedule. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision to test for TRAIL-R antibodies by dot blot assay, ELISA (Enzyme-Linked Immunosorbent Assay) or inhibition of TRAIL binding.

Following detection of an appropriate antibody titer, positive animals are provided one last intravenous injection of TRAIL-R in saline. Three to four days later, the animals are sacrificed, spleen cells harvested, and spleen cells are fused to a murine myeloma cell line, e.g., NS1 or preferably P3x63Ag8.653 (ATCC CRL 1580). Fusions generate hybridoma cells, which are plated in multiple microtiter plates in a HAT (hypoxanthine, aminopterin and thymidine) selective medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells are screened by ELISA for reactivity against purified TRAIL-R by adaptations of the techniques disclosed in Engvall et al., *Immunochem.* 8:871, 1971 and in U.S. Patent 4,703,004. A preferred screening technique is the antibody capture technique described in Beckmann et al., (*J. Immunol.* 144:4212, 1990). Positive hybridoma cells can be injected intraperitoneally into syngeneic BALB/c mice to produce ascites containing high concentrations of anti-TRAIL-R monoclonal antibodies. Alternatively, hybridoma cells can be grown *in vitro* in flasks or roller bottles by various techniques. Monoclonal antibodies produced in mouse ascites can be purified by ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to Protein A or Protein G can also be used, as can affinity chromatography based upon binding to TRAIL-R.

#### **EXAMPLE 5: Northern Blot Analysis**

The tissue distribution of TRAIL-R mRNA was investigated by Northern blot analysis, as follows. An aliquot of a radiolabeled probe (the same radiolabeled probe used to screen the cDNA library in example 3) was added to two different human multiple tissue Northern blots (Clontech, Palo Alto, CA; Biochain, Palo Alto, CA). Hybridization was conducted overnight at 63°C in 50% formamide as previously described (March et al., *Nature* 315:641-647, 1985). The blots then were washed with 2X SSC, 0.1% SDS at 68°C for 30 minutes. A glycerol-aldehyde-phosphate dehydrogenase (GAPDH) specific probe was used to standardize for RNA loadings.

A single transcript of 4.4 kilobases (kb) was present in all tissues examined, including spleen, thymus, peripheral blood lymphocytes (PBLs), prostate, testis, ovary,



uterus, placenta, and multiple tissues along the gastro-intestinal tract (including esophagus, stomach, duodenum, jejunum/ileum, colon, rectum, and small intestine). The cells and tissues with the highest levels of TRAIL-R mRNA are PBLs, spleen, and ovary, as shown by comparison to control hybridizations with a GAPDH-specific probe.

5

#### **EXAMPLE 6: Binding Assay**

Full length human TRAIL-R was expressed and tested for the ability to bind TRAIL. The binding assay was conducted as follows.

10 A fusion protein comprising a leucine zipper peptide fused to the N-terminus of a soluble TRAIL polypeptide (LZ-TRAIL) was employed in the assay. An expression construct was prepared, essentially as described for preparation of the Flag®-TRAIL expression construct in Wiley et al. (*Immunity*, 3:673-682, 1995; hereby incorporated by reference), except that DNA encoding the Flag® peptide was replaced with a sequence encoding a modified leucine zipper that allows for trimerization. The construct, in  
15 expression vector pDC409, encoded a leader sequence derived from human cytomegalovirus, followed by the leucine zipper moiety fused to the N-terminus of a soluble TRAIL polypeptide. The TRAIL polypeptide comprised amino acids 95-281 of human TRAIL (a fragment of the extracellular domain), as described in Wiley et al. (*supra*). The LZ-TRAIL was expressed in CHO cells, and purified from the culture  
20 supernatant.

The expression vector designated pDC409 is a mammalian expression vector derived from the pDC406 vector described in McMahan et al. (*EMBO J.* 10:2821-2832, 1991; hereby incorporated by reference). Features added to pDC409 (compared to pDC406) include additional unique restriction sites in the multiple cloning site (mcs); three  
25 stop codons (one in each reading frame) positioned downstream of the mcs; and a T7 polymerase promoter, downstream of the mcs, that facilitates sequencing of DNA inserted into the mcs.

For expression of full length human TRAIL-R protein, the entire coding region (i.e., the DNA sequence presented in SEQ ID NO:1) was amplified by polymerase chain  
30 reaction (PCR). The template employed in the PCR was the cDNA clone isolated from a human foreskin fibroblast cDNA library, as described in example 3. The isolated and amplified DNA was inserted into the expression vector pDC409, to yield a construct designated pDC409-TRAIL-R.

35 CrmA protein was employed to inhibit apoptosis of host cells expressing recombinant TRAIL-R, as discussed above and in example 8. An expression vector designated pDC409-CrmA contained DNA encoding poxvirus CrmA in pDC409. The 38-

kilodalton cowpox-derived protein that was subsequently designated CrmA is described in Pickup et al. (*Proc. Natl. Acad. Sci. USA* 83:7698-7702, 1986; hereby incorporated by reference).

CV-1/EBNA cells were co-transfected with pDC409-TRAIL-R together with  
5 pDC409-CrmA, or with pDC409-CrmA alone. The cells were cultured in DMEM supplemented with 10% fetal bovine serum, penicillin, streptomycin, and glutamine. 48 hours after transfection, cells were detached non-enzymatically and incubated with LZ-TRAIL (5 µg/ml), a biotinylated anti-LZ monoclonal antibody (5 µg/ml), and phycoerythrin-conjugated streptavidin (1:400), before analysis by fluorescence-activated  
10 cell scanning (FACS). The cytometric analysis was conducted on a FACScan (Beckton Dickinson, San Jose, CA).

The CV-1/EBNA cells co-transfected with vectors encoding TRAIL-R and CrmA showed significantly enhanced binding of LZ-TRAIL, compared to the cells transfected with the CrmA-encoding vector alone.

15

**EXAMPLE 7: TRAIL-R Blocks TRAIL-Induced Apoptosis**  
**of Target Cells**

TRAIL-R was tested for the ability to block TRAIL-induced apoptosis of Jurkat cells. The TRAIL-R employed in the assay was in the form of a fusion protein designated  
20 sTRAIL-R/Fc, which comprised the extracellular domain of human TRAIL-R, fused to the N-terminus of an Fc polypeptide derived from human IgG1.

CV1-EBNA cells were transfected with a recombinant expression vector comprising a gene fusion encoding the sTRAIL-R/Fc protein, in the pDC409 vector described in example 6, and cultured to allow expression of the fusion protein. The  
25 sTRAIL-R/Fc fusion protein was recovered from the culture supernatant. Procedures for fusing DNA encoding an IgG1 Fc polypeptide to DNA encoding a heterologous protein are described in Smith et al., (*Cell* 73:1349-1360, 1993); analogous procedures were employed herein.

A fusion protein designated TNF-R2-Fc, employed as a control, comprised the  
30 extracellular domain of the TNF receptor protein known as p75 or p80 TNF-R (Smith et al., *Science* 248:1019-1023, 1990; Smith et al. *Cell* 76:959-962, 1994), fused to an Fc polypeptide. A mouse monoclonal antibody that is a blocking antibody directed against human TRAIL, was employed in the assay as well.

Jurkat cells were incubated with varying or constant concentrations of LZ-TRAIL  
35 (the LZ-TRAIL fusion protein described in example 6), in the absence or presence of varying concentrations of sTRAIL-R-Fc, TNF-R2-Fc, or the TRAIL-specific monoclonal

antibody. Cell death was quantitated using an MTT cell viability assay (MTT is 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide), as previously described (Mosmann, *J. Immunol. Methods* 65:55-63, 1983). The results are shown in Figure 2, which presents the percent cell death for Jurkat cells that were untreated ( $\Delta$ ) or were treated with varying ( $\blacktriangle$ ) or constant ( $\circ$ ,  $\bullet$ ,  $\square$ ,  $\blacksquare$ ) concentrations of LZ-TRAIL (13 ng/ml) in the absence ( $\bullet$ ) or presence of varying concentrations of TRAIL-R2-Fc ( $\blacksquare$ ), TNF-R2-Fc ( $\square$ ), or the anti-TRAIL antibody ( $\circ$ ). Varying concentrations for all substances started at 10  $\mu$ g/ml and were serially diluted.

The anti-TRAIL monoclonal antibody and sTRAIL-R/Fc each blocked TRAIL-induced apoptosis in a dose dependent fashion, whereas TNFR2-Fc did not. Thus, the extracellular domain of TRAIL-R is capable of binding to TRAIL and inhibiting TRAIL-mediated apoptosis of target cells.

#### **EXAMPLE 8: TRAIL-R-induced apoptosis is blocked by caspase inhibitors and FADD-DN**

CV-1/EBNA cells were transfected, by the DEAE-dextran method, with expression plasmids for TRAIL-R (pDC409-TRAIL-R), together with a threefold excess of empty expression vector (pDC409) in the presence or absence of z-VAD-fmk (10 $\mu$ M; in the culture medium), or together with a threefold excess of expression vector pDC409-CrmA, pDC409-p35, or pDC409-FADD-DN. In addition, 400 ng/slide of an expression vector for the *E. coli lacz* gene was co-transfected together with all DNA mixes. The transfected cells were cultured in chambers mounted on slides.

The mammalian expression vector pDC409, and the pDC409-TRAIL-R vector encoding full length human TRAIL-R, are described in example 6. The tripeptide derivative zVAD-fmk (benzyloxy-carbonyl-Val-Ala-Asp-fluoromethylketone) is available from Enzyme System Products, Dublin, California.

The 38-kilodalton cowpox-derived protein that was subsequently designated CrmA is described in Pickup et al. (*Proc. Natl. Acad. Sci. USA* 83:7698-7702, 1986; hereby incorporated by reference). Sequence information for CrmA is presented in Figure 4 of Pickup et al., *supra*.

A 35-kilodalton protein encoded by *Autographa californica* nuclear polyhedrosis virus, a baculovirus, is described in Friesen and Miller (*J. Virol.* 61:2264-2272, 1987; hereby incorporated by reference). Sequence information for this protein, designated baculovirus p35 herein, is presented in Figure 5 of Friesen and Miller, *supra*.

FADD (also designated MORT1) is described in Boldin et al. (*J. Biol. Chem.* 270:7795-7798, 1995; hereby incorporated by reference). The protein referred to as

FADD-DN (FADD dominant negative) is a C-terminal fragment of FADD that includes the death domain. DNA encoding FADD-DN, fused to an N-terminal Flag<sup>®</sup> epitope tag (described above), was inserted into the pDC409 expression vector described in example 6, to form pDC409-FADD-DN. The FADD-DN polypeptide corresponds to amino acids 117 through 245 of the MORT1 amino acid sequence presented in Boldin et al., *supra*.

48 hours after transfection, cells were washed with PBS, fixed with glutaraldehyde and incubated with X-gal (5-bromo-4-chloro-3-indoxyl- $\beta$ -D-galactopyranoside). Cells expressing  $\beta$ -galactosidase stain blue. A decrease in the percentage of stained cells indicates loss of  $\beta$ -galactosidase expression, and correlates with death of cells that express the protein(s) co-transfected with the *lacZ* gene.

The results are presented in Figure 3, wherein the values plotted represent the mean and standard deviation of at least three separate experiments. Poxvirus CrmA, baculovirus p35, FADD-DN, and z-VAD-fmk each effectively reduced death of transfected cells expressing TRAIL-R.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(i) APPLICANT: Rauch, Charles  
Walczak, Henning

(ii) TITLE OF INVENTION: Receptor That Binds TRAIL

(iii) NUMBER OF SEQUENCES: 4

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Kathryn A. Anderson, Immunex Corporation  
(B) STREET: 51 University Street  
(C) CITY: Seattle,  
(D) STATE: WA  
(E) COUNTRY: US  
(F) ZIP: 98101

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC Compatible  
(C) OPERATING SYSTEM: MS-DOS/Windows 95  
(D) SOFTWARE: Word for Windows 95, 7.0a

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: --to be assigned--  
(B) FILING DATE: 11-FEB-1998  
(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/883,036  
(B) FILING DATE: 26-JUN-1997

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/869,852  
(B) FILING DATE: 04-JUN-1997

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/829,536  
(B) FILING DATE: 28-MAR-1997

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/815,255  
(B) FILING DATE: 12-MAR-1997

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/799,861  
(B) FILING DATE: 13-FEB-1997

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Anderson, Kathryn A.  
(B) REGISTRATION NUMBER: 32,172  
(C) REFERENCE/DOCKET NUMBER: 2625-WO

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 587-0430  
 (B) TELEFAX: (206) 233-0644  
 (C) TELEX: 756822

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1323 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vii) IMMEDIATE SOURCE:

(B) CLONE: huTrail-R

## (ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 1..1323

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG	GAA	CAA	CGG	GGA	CAG	AAC	GCC	CCG	GCC	GCT	TCG	GGG	GCC	CGG	AAA	48
Met	Glu	Gln	Arg	Gly	Gln	Asn	Ala	Pro	Ala	Ala	Ser	Gly	Ala	Arg	Lys	
1				5					10					15		
AGG	CAC	GGC	CCA	GGA	CCC	AGG	GAG	GCG	CGG	GGA	GCC	AGG	CCT	GGG	CCC	96
Arg	His	Gly	Pro	Gly	Pro	Arg	Glu	Ala	Arg	Gly	Ala	Arg	Pro	Gly	Pro	
			20					25					30			
CGG	GTC	CCC	AAG	ACC	CTT	GTG	CTC	GTT	GTC	GCC	GCG	GTC	CTG	CTG	TTG	144
Arg	Val	Pro	Lys	Thr	Leu	Val	Leu	Val	Val	Ala	Ala	Val	Leu	Leu	Leu	
		35				40						45				
GTC	TCA	GCT	GAG	TCT	GCT	CTG	ATC	ACC	CAA	CAA	GAC	CTA	GCT	CCC	CAG	192
Val	Ser	Ala	Glu	Ser	Ala	Leu	Ile	Thr	Gln	Gln	Asp	Leu	Ala	Pro	Gln	
	50					55					60					
CAG	AGA	GCG	GCC	CCA	CAA	CAA	AAG	AGG	TCC	AGC	CCC	TCA	GAG	GGA	TTG	240
Gln	Arg	Ala	Ala	Pro	Gln	Gln	Lys	Arg	Ser	Ser	Pro	Ser	Glu	Gly	Leu	
65				70					75					80		
TGT	CCA	CCT	GGA	CAC	CAT	ATC	TCA	GAA	GAC	GGT	AGA	GAT	TGC	ATC	TCC	288
Cys	Pro	Pro	Gly	His	His	Ile	Ser	Glu	Asp	Gly	Arg	Asp	Cys	Ile	Ser	
			85						90				95			
TGC	AAA	TAT	GGA	CAG	GAC	TAT	AGC	ACT	CAC	TGG	AAT	GAC	CTC	CTT	TTC	336
Cys	Lys	Tyr	Gly	Gln	Asp	Tyr	Ser	Thr	His	Trp	Asn	Asp	Leu	Leu	Phe	
			100					105					110			

TGC	TTG	CGC	TGC	ACC	AGG	TGT	GAT	TCA	GGT	GAA	GTG	GAG	CTA	AGT	CCG	384
Cys	Leu	Arg	Cys	Thr	Arg	Cys	Asp	Ser	Gly	Glu	Val	Glu	Leu	Ser	Pro	
		115					120					125				
TGC	ACC	ACG	ACC	AGA	AAC	ACA	GTG	TGT	CAG	TGC	GAA	GAA	GGC	ACC	TTC	432
Cys	Thr	Thr	Thr	Arg	Asn	Thr	Val	Cys	Gln	Cys	Glu	Glu	Gly	Thr	Phe	
	130					135					140					
CGG	GAA	GAA	GAT	TCT	CCT	GAG	ATG	TGC	CGG	AAG	TGC	CGC	ACA	GGG	TGT	480
Arg	Glu	Glu	Asp	Ser	Pro	Glu	Met	Cys	Arg	Lys	Cys	Arg	Thr	Gly	Cys	
145					150					155					160	
CCC	AGA	GGG	ATG	GTC	AAG	GTC	GGT	GAT	TGT	ACA	CCC	TGG	AGT	GAC	ATC	528
Pro	Arg	Gly	Met	Val	Lys	Val	Gly	Asp	Cys	Thr	Pro	Trp	Ser	Asp	Ile	
				165					170					175		
GAA	TGT	GTC	CAC	AAA	GAA	TCA	GGT	ACA	AAG	CAC	AGT	GGG	GAA	GCC	CCA	576
Glu	Cys	Val	His	Lys	Glu	Ser	Gly	Thr	Lys	His	Ser	Gly	Glu	Ala	Pro	
			180					185					190			
GCT	GTG	GAG	GAG	ACG	GTG	ACC	TCC	AGC	CCA	GGG	ACT	CCT	GCC	TCT	CCC	624
Ala	Val	Glu	Glu	Thr	Val	Thr	Ser	Ser	Pro	Gly	Thr	Pro	Ala	Ser	Pro	
		195					200					205				
TGT	TCT	CTC	TCA	GGC	ATC	ATC	ATA	GGA	GTC	ACA	GTT	GCA	GCC	GTA	GTC	672
Cys	Ser	Leu	Ser	Gly	Ile	Ile	Ile	Gly	Val	Thr	Val	Ala	Ala	Val	Val	
	210					215					220					
TTG	ATT	GTG	GCT	GTG	TTT	GTT	TGC	AAG	TCT	TTA	CTG	TGG	AAG	AAA	GTC	720
Leu	Ile	Val	Ala	Val	Phe	Val	Cys	Lys	Ser	Leu	Leu	Trp	Lys	Lys	Val	
225					230					235					240	
CTT	CCT	TAC	CTG	AAA	GGC	ATC	TGC	TCA	GGT	GGT	GGT	GGG	GAC	CCT	GAG	768
Leu	Pro	Tyr	Leu	Lys	Gly	Ile	Cys	Ser	Gly	Gly	Gly	Gly	Asp	Pro	Glu	
				245					250					255		
CGT	GTG	GAC	AGA	AGC	TCA	CAA	CGA	CCT	GGG	GCT	GAG	GAC	AAT	GTC	CTC	816
Arg	Val	Asp	Arg	Ser	Ser	Gln	Arg	Pro	Gly	Ala	Glu	Asp	Asn	Val	Leu	
			260					265					270			
AAT	GAG	ATC	GTG	AGT	ATC	TTG	CAG	CCC	ACC	CAG	GTC	CCT	GAG	CAG	GAA	864
Asn	Glu	Ile	Val	Ser	Ile	Leu	Gln	Pro	Thr	Gln	Val	Pro	Glu	Gln	Glu	
		275					280					285				
ATG	GAA	GTC	CAG	GAG	CCA	GCA	GAG	CCA	ACA	GGT	GTC	AAC	ATG	TTG	TCC	912
Met	Glu	Val	Gln	Glu	Pro	Ala	Glu	Pro	Thr	Gly	Val	Asn	Met	Leu	Ser	
	290					295					300					
CCC	GGG	GAG	TCA	GAG	CAT	CTG	CTG	GAA	CCG	GCA	GAA	GCT	GAA	AGG	TCT	960
Pro	Gly	Glu	Ser	Glu	His	Leu	Leu	Glu	Pro	Ala	Glu	Ala	Glu	Arg	Ser	
305					310					315					320	
CAG	AGG	AGG	AGG	CTG	CTG	GTT	CCA	GCA	AAT	GAA	GGT	GAT	CCC	ACT	GAG	1008
Gln	Arg	Arg	Arg	Leu	Leu	Val	Pro	Ala	Asn	Glu	Gly	Asp	Pro	Thr	Glu	
				325					330					335		
ACT	CTG	AGA	CAG	TGC	TTC	GAT	GAC	TTT	GCA	GAC	TTG	GTG	CCC	TTT	GAC	1056
Thr	Leu	Arg	Gln	Cys	Phe	Asp	Asp	Phe	Ala	Asp	Leu	Val	Pro	Phe	Asp	
			340					345					350			

TCC	TGG	GAG	CCG	CTC	ATG	AGG	AAG	TTG	GGC	CTC	ATG	GAC	AAT	GAG	ATA	1104
Ser	Trp	Glu	Pro	Leu	Met	Arg	Lys	Leu	Gly	Leu	Met	Asp	Asn	Glu	Ile	
		355					360					365				
AAG	GTG	GCT	AAA	GCT	GAG	GCA	GCG	GGC	CAC	AGG	GAC	ACC	TTG	TAC	ACG	1152
Lys	Val	Ala	Lys	Ala	Glu	Ala	Ala	Gly	His	Arg	Asp	Thr	Leu	Tyr	Thr	
	370					375					380					
ATG	CTG	ATA	AAG	TGG	GTC	AAC	AAA	ACC	GGG	CGA	GAT	GCC	TCT	GTC	CAC	1200
Met	Leu	Ile	Lys	Trp	Val	Asn	Lys	Thr	Gly	Arg	Asp	Ala	Ser	Val	His	
385					390					395					400	
ACC	CTG	CTG	GAT	GCC	TTG	GAG	ACG	CTG	GGA	GAG	AGA	CTT	GCC	AAG	CAG	1248
Thr	Leu	Leu	Asp	Ala	Leu	Glu	Thr	Leu	Gly	Glu	Arg	Leu	Ala	Lys	Gln	
				405					410					415		
AAG	ATT	GAG	GAC	CAC	TTG	TTG	AGC	TCT	GGA	AAG	TTC	ATG	TAT	CTA	GAA	1296
Lys	Ile	Glu	Asp	His	Leu	Leu	Ser	Ser	Gly	Lys	Phe	Met	Tyr	Leu	Glu	
			420					425					430			
GGT	AAT	GCA	GAC	TCT	GCC	ATG	TCC	TAA								1323
Gly	Asn	Ala	Asp	Ser	Ala	Met	Ser	*								
		435				440										

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 440 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Glu	Gln	Arg	Gly	Gln	Asn	Ala	Pro	Ala	Ala	Ser	Gly	Ala	Arg	Lys
1				5					10					15	
Arg	His	Gly	Pro	Gly	Pro	Arg	Glu	Ala	Arg	Gly	Ala	Arg	Pro	Gly	Pro
			20					25					30		
Arg	Val	Pro	Lys	Thr	Leu	Val	Leu	Val	Val	Ala	Ala	Val	Leu	Leu	Leu
	35					40						45			
Val	Ser	Ala	Glu	Ser	Ala	Leu	Ile	Thr	Gln	Gln	Asp	Leu	Ala	Pro	Gln
	50					55					60				
Gln	Arg	Ala	Ala	Pro	Gln	Gln	Lys	Arg	Ser	Ser	Pro	Ser	Glu	Gly	Leu
	65				70				75					80	
Cys	Pro	Pro	Gly	His	His	Ile	Ser	Glu	Asp	Gly	Arg	Asp	Cys	Ile	Ser
				85					90				95		
Cys	Lys	Tyr	Gly	Gln	Asp	Tyr	Ser	Thr	His	Trp	Asn	Asp	Leu	Leu	Phe
			100					105					110		



Cys Leu Arg Cys Thr Arg Cys Asp Ser Gly Glu Val Glu Leu Ser Pro  
 115 120 125  
 Cys Thr Thr Thr Arg Asn Thr Val Cys Gln Cys Glu Glu Gly Thr Phe  
 130 135 140  
 Arg Glu Glu Asp Ser Pro Glu Met Cys Arg Lys Cys Arg Thr Gly Cys  
 145 150 155 160  
 Pro Arg Gly Met Val Lys Val Gly Asp Cys Thr Pro Trp Ser Asp Ile  
 165 170 175  
 Glu Cys Val His Lys Glu Ser Gly Thr Lys His Ser Gly Glu Ala Pro  
 180 185 190  
 Ala Val Glu Glu Thr Val Thr Ser Ser Pro Gly Thr Pro Ala Ser Pro  
 195 200 205  
 Cys Ser Leu Ser Gly Ile Ile Ile Gly Val Thr Val Ala Ala Val Val  
 210 215 220  
 Leu Ile Val Ala Val Phe Val Cys Lys Ser Leu Leu Trp Lys Lys Val  
 225 230 235 240  
 Leu Pro Tyr Leu Lys Gly Ile Cys Ser Gly Gly Gly Gly Asp Pro Glu  
 245 250 255  
 Arg Val Asp Arg Ser Ser Gln Arg Pro Gly Ala Glu Asp Asn Val Leu  
 260 265 270  
 Asn Glu Ile Val Ser Ile Leu Gln Pro Thr Gln Val Pro Glu Gln Glu  
 275 280 285  
 Met Glu Val Gln Glu Pro Ala Glu Pro Thr Gly Val Asn Met Leu Ser  
 290 295 300  
 Pro Gly Glu Ser Glu His Leu Leu Glu Pro Ala Glu Ala Glu Arg Ser  
 305 310 315 320  
 Gln Arg Arg Arg Leu Leu Val Pro Ala Asn Glu Gly Asp Pro Thr Glu  
 325 330 335  
 Thr Leu Arg Gln Cys Phe Asp Asp Phe Ala Asp Leu Val Pro Phe Asp  
 340 345 350  
 Ser Trp Glu Pro Leu Met Arg Lys Leu Gly Leu Met Asp Asn Glu Ile  
 355 360 365  
 Lys Val Ala Lys Ala Glu Ala Ala Gly His Arg Asp Thr Leu Tyr Thr  
 370 375 380  
 Met Leu Ile Lys Trp Val Asn Lys Thr Gly Arg Asp Ala Ser Val His  
 385 390 395 400  
 Thr Leu Leu Asp Ala Leu Glu Thr Leu Gly Glu Arg Leu Ala Lys Gln  
 405 410 415  
 Lys Ile Glu Asp His Leu Leu Ser Ser Gly Lys Phe Met Tyr Leu Glu  
 420 425 430

Gly Asn Ala Asp Ser Ala Met Ser \*  
 435 440

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 157 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: huTrail-R frag

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 3..155

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CT GAG ACT CTG AGA CAG TGC TTC GAT GAC TTT GCA GAC TTG GTG CCC	47
Glu Thr Leu Arg Gln Cys Phe Asp Asp Phe Ala Asp Leu Val Pro	
1 5 10 15	
TTT GAC TCC TGG GAG CCG CTC ATG AGG AAG TTG GGC CTC ATG GAC AAT	95
Phe Asp Ser Trp Glu Pro Leu Met Arg Lys Leu Gly Leu Met Asp Asn	
20 25 30	
GAG ATA AAG GTG GCT AAA GCT GAG GCA GCG GGC CAC AGG GAC ACC TTG	143
Glu Ile Lys Val Ala Lys Ala Glu Ala Ala Gly His Arg Asp Thr Leu	
35 40 45	
TNC ACN ATG CTG AT	157
Xaa Thr Met Leu	
50	

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 51 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

(2) INFORMATION FOR SEQ ID NO:5:

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:  
(B) CLONE: FLAG peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Tyr Lys Asp Asp Asp Asp Lys  
1 5

What is claimed is:

1. A purified TRAIL receptor (TRAIL-R) polypeptide that is capable of binding TRAIL, wherein the TRAIL-R is characterized as comprising the amino acid sequence VPANEGD.
2. A TRAIL-R polypeptide of claim 1, wherein said polypeptide is further characterized by a molecular weight of about 50 to 55 kilodaltons.
3. A TRAIL-R polypeptide of claim 1, wherein said polypeptide is further characterized by comprising the amino acid sequence ETLRQCFDDFADLVPFDS WEPLMRKLGLMDNEIKVAKAEAAGHRDTLXTML.
4. A TRAIL-R polypeptide of claim 3, wherein said polypeptide is further characterized by a molecular weight of about 50 to 55 kilodaltons.
5. A purified TRAIL-R polypeptide selected from the group consisting of:
  - a) the TRAIL-R polypeptide of SEQ ID NO:2; and
  - b) a fragment of the polypeptide of (a), wherein said fragment is capable of binding TRAIL.
6. A TRAIL-R polypeptide of claim 5, wherein said polypeptide comprises amino acids x to 440 of SEQ ID NO:2, wherein x represents an integer from 51 through 59.
7. A TRAIL-R polypeptide of claim 6, wherein said polypeptide comprises amino acids 54 to 440 of SEQ ID NO:2.
8. A TRAIL-R polypeptide of claim 5, wherein said fragment is a soluble TRAIL-R comprising the extracellular domain of the TRAIL-R protein of SEQ ID NO:2.
9. A purified TRAIL-R polypeptide comprising an amino acid sequence that is at least 80% identical to the amino acid sequence presented in SEQ ID NO:2.

10. A TRAIL-R polypeptide of claim 9, wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to the amino acid sequence presented in SEQ ID NO:2.

11. A TRAIL-R polypeptide of claim 10, wherein said polypeptide comprises an amino acid sequence that is at least 95% identical to the amino acid sequence presented in SEQ ID NO:2.

12. A TRAIL-R polypeptide of claim 9, wherein said polypeptide is naturally occurring.

13. An oligomer comprising from two to four TRAIL-R polypeptides of claim 5.

14. An oligomer comprising from two to four TRAIL-R polypeptides of claim 8.

15. A composition comprising a TRAIL-R polypeptide of claim 5, and a physiologically acceptable diluent, excipient, or carrier.

16. A composition comprising a TRAIL-R polypeptide of claim 8, and a physiologically acceptable diluent, excipient, or carrier.

17. An isolated TRAIL-R DNA, wherein said DNA comprises the nucleotide sequence presented in Figure 1.

18. An isolated TRAIL-R DNA, wherein said DNA encodes a polypeptide selected from the group consisting of:

- a) the TRAIL-R polypeptide of SEQ ID NO:2; and
- b) a fragment of the polypeptide of (a), wherein said fragment is capable of binding TRAIL.

19. A TRAIL-R DNA of claim 18, wherein said DNA encodes amino acids 1 to 440 of SEQ ID NO:2.

20. A TRAIL-R DNA of claim 18, wherein said polypeptide comprises amino acids x to 440 of SEQ ID NO:2, wherein x represents an integer from 51 through 59.

21. A TRAIL-R DNA of claim 20, wherein said polypeptide comprises amino acids 54 to 440 of SEQ ID NO:2.

22. A TRAIL-R DNA of claim 18, wherein said fragment is a soluble TRAIL-R comprising the extracellular domain of the TRAIL-R protein of SEQ ID NO:2.

23. An isolated TRAIL-R DNA, wherein said DNA encodes a polypeptide comprising an amino acid sequence that is at least 80% identical to the amino acid sequence presented in SEQ ID NO:2.

24. A TRAIL-R DNA of claim 23, wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to the amino acid sequence presented in SEQ ID NO:2.

25. A TRAIL-R DNA of claim 24, wherein said polypeptide comprises an amino acid sequence that is at least 95% identical to the amino acid sequence presented in SEQ ID NO:2.

26. A TRAIL-R DNA of claim 23, wherein said polypeptide is naturally occurring.

27. An expression vector comprising a DNA according to claim 18.

28. An expression vector comprising a DNA according to claim 19.

29. An expression vector comprising a DNA according to claim 20.

30. An expression vector comprising a DNA according to claim 22.

31. An expression vector comprising a DNA according to claim 23.

32. A host cell transformed with an expression vector of claim 27.

33. A host cell transformed with an expression vector of claim 28.
34. A host cell transformed with an expression vector of claim 29.
35. A host cell transformed with an expression vector of claim 30.
36. A host cell transformed with an expression vector of claim 31.
37. An isolated TRAIL-R DNA comprising at least 60 nucleotides of the sequence of SEQ ID NO:1, or the DNA or RNA complement thereof.
38. An antibody that is directed against a TRAIL-R polypeptide of claim 5, or an antigen-binding fragment of said antibody.
39. An antibody of claim 38, wherein the antibody is a monoclonal antibody.

## FIGURE 1

1 CTGAGACTCTGAGACAGTGCTTCGATGACTTTGCAGACTT 40  
-----+-----+-----+-----+  
GACTCTGAGACTCTGTCACGAAGCTACTGAAACGTCTGAA  
E T L R Q C F D D F A D L

41 GGTGCCCTTTGACTCCTGGGAGCCGCTCATGAGGAAGTTG 80  
-----+-----+-----+-----+  
CCACGGGAAACTGAGGACCCTCGGCGAGTACTCCTTCAAC  
V P F D S W E P L M R K L

81 GGCCTCATGGACAATGAGATAAAGGTGGCTAAAGCTGAGG 120  
-----+-----+-----+-----+  
CCGGAGTACCTGTTACTCTATTTCCACCGATTTCGACTCC  
G L M D N E I K V A K A E A

121 CAGCGGGCCACAGGGACACCTTGTNCACNATGCTGAT 157  
-----+-----+-----+-----+  
GTCGCCCCGGTGTCCCTGTGGAACANGTGNTACGACTA  
A G H R D T L X T M L



FIGURE 2





FIGURE 3



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/02239

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) :C07K 14/435, 14/715, 16/28; C12N 15/12, 15/19 US CL :530/350, 351, 387.9; 536/23.5 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 530/350, 351, 387.9; 536/23.5  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, CAPLUS, MEDLINE, BIOSIS, SCISEARCH search terms: trail, apoptosis, receptor, antibody, DNA, nucleic acid, clone, cDNA		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	PAN, et al. The receptor for cytotoxic ligand TRAIL. 04 April 1997. Vol 276, No. 5309, pages 111-113, see entire document.	1-39
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* *A* *E* *L* *O* *P*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *A* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 15 MAY 1998		Date of mailing of the international search report 19 JUN 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer DARYL A. BASHAM Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/02239

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/02239

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-37, drawn to polypeptide, an oligomer, a composition, and isolated DNA, an expression vector and a host cell.

Group II, claim(s) 38 and 39, drawn to an antibody.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The polypeptide and DNA compositions of Group I have materially different chemical structures and biological functions from the antibodies of Group II. The special technical feature by which the polypeptide and DNA of Group I are defined distinguish them from the special technical feature which defines the antibody of Group II and as they do not share identical structural and functional features, including that they are distinct compositions, they do not share the same or corresponding special technical features and unity of invention is lacking. The claims are not so linked by a special technical feature within the meaning of the PCT Rule 13.2 so as to for a single inventive concept.